Transiting Exoplanet Survey Satellite (TESS) Observations of Pulsations in Eclipsing Binary Systems Containing Delta Scuti Pulsating Components

Argent Astro Research Institute (AARI)

Abstract: The Transiting Exoplanet Survey Satellite (TESS) observed pulsations in 34 eclipsing binary systems containing Delta Scuti pulsating components in the instability strip of the Pleiades open cluster. TESS is producing high-precision, rapid-cadence light curves over most of the sky, opening up new possibilities for studying large samples of Delta Scuti stars, both singly and in binary systems. Using time-series photometry data from TESS contained in the Mikulski Archive for Space Telescopes (MAST) database, light curves and periodograms are examined in detail to extract frequency/period relations among the pulsations as well as the orbital periods of the eclipsing binary pairs.

Introduction and Background

Bedding, T. R., et al., 2023, studied 89 A- and F-type members of the Pleiades open cluster. Using time-series photometry from the NASA Transiting Exoplanet Survey Satellite (TESS), they discovered Delta Scuti (δ-Scuti) pulsations in 34 stars. Additionally, Hinzel, D.H., 2025, studied TESS observations of eclipsing binary V477 Cygni which included a δ-Scuti pulsator as one of its components. TESS is producing high-precision, rapid-cadence light curves over most of the sky, opening up new possibilities for studying large samples of δ-Scuti stars. TESS observations are made in 27 day sectors. The observations of the 34 systems in the Pleiades cluster were made in the fourth year of the mission in Sectors 42-44 (August 20 to November 6, 2021). The TESS observations incorporate a 120 second cadence for all objects. All 89 members of the Pleiades open cluster, including the 34 δ-Scuti pulsators, are summarized in Table 1 of Bedding, T. R., et al., 2023. All objects are assigned a TESS Input Catalog (TIC) designation number, but Table 1 also lists common names for most stars such as HD 23156, V624 Tau, etc. For each system a light curve and periodogram of the orbital period and pulsation spectrum are presented.

Methods

This study looks at observed pulsations in 34 eclipsing binary systems containing δ-Scuti pulsating components. The light curve and the periodograms are analyzed from TESS photometry data contained in the Mikulski Archive for Space Telescopes (MAST). All of the TESS photometry files were downloaded and a light curve was generated by the American Association of Variable Star Observers (AAVSO) VStar software (Benn, D. 2012, Algorithms + Observations = VStar, Journal of the AAVSO (JAAVSO), v40, n2, pp.852-866). Prior to mode identification of pulsation mode frequencies (or periods), the frequencies themselves must be extracted by assembling time-series data of the star's

observable surface properties. Such time-series data can be obtained from time-series photometry with the periodic variability of the star's brightness as a function of time. This is the star's light curve. To extract pulsation mode frequencies, it is common practice to employ Fourier analysis, specifically Discrete Fourier Transforms (DFTs) for unevenly sampled time series or a Lomb-Scargle periodogram (Scargle, 1982). The frequency spectrum is calculated up to the Nyquist frequency, defined as: F (Nyquist) = 1/2Δt where Δt is the cadence of the time series. Once the frequency spectrum has been calculated, peaks are noted that are statistically significant and represent pulsation mode frequencies based on satisfying a significance criterion. This is the periodogram. The data is analyzed with VStar. Specifically, Fourier analysis of the combined photometric data is performed to yield a detailed periodogram for the δ-Scuti pulsator from which frequencies, periodicities, and other variations can potentially be identified. VStar utilizes the Date Compensated Discrete Fourier Transform (DCDFT) algorithm (Ferraz-Mello, 1981) to produce a power spectrum, a period range, and a resolution. The Date Compensated DFT compensates for gaps in the data, which is common for variable star observations. The resulting analysis can include one or more periods, one or more harmonics, and one or more subharmonics. These can be selected to create a model that can also include a polynomial function that is used as a smoothing mechanism to capture key aspects of the data set without all the noise and fine fluctuations. When a model is created, it is subtracted from observations in the series to yield a second series called a residual. The residuals can also be analyzed to look for other signals (frequencies/periods) in a process called pre-whitening. Periodicities and other potential variations are analyzed utilizing TESS photometry, models are created from the photometry, mean series are computed, and residuals are analyzed to obtain all possible variations.

Analysis and Results

Presented below are the light curves, orbital period periodograms, and pulsation spectra periodograms for 34 eclipsing binary systems containing δ-Scuti pulsators.



Light Curve for TIC 440691730



Period and Pulsation Analysis (DCDFT) for TIC 440691730



Light Curve for TIC 440681425



Period and Pulsation Analysis (DCDFT) for TIC 440681425



Light Curve for TIC 405483425



Period and Pulsation Analysis (DCDFT) for TIC 405483425



Light Curve for TIC 402366726



Period and Pulsation Analysis (DCDFT) for TIC 402366726



Light Curve for TIC 385554826



Period and Pulsation Analysis (DCDFT) for TIC 385554826



Light Curve for TIC 346626099



Period and Pulsation Analysis (DCDFT) for TIC 346626099



Light Curve for TIC 346626001



Period and Pulsation Analysis (DCDFT) for TIC 346626001



Light Curve for TIC 125736946



Period and Pulsation Analysis (DCDFT) for TIC 125736946



Light Curve for TIC 125736216



Period and Pulsation Analysis (DCDFT) for TIC 125736216



Light Curve for TIC 67828699



Period and Pulsation Analysis (DCDFT) for TIC 67828699



Light Curve for TIC 29058513



Period and Pulsation Analysis (DCDFT) for TIC 29058513



Light Curve for TIC 46476992



Period and Pulsation Analysis (DCDFT) for TIC 46476992



Light Curve for TIC 67830321



Period and Pulsation Analysis (DCDFT) for TIC 67830321



Light Curve for TIC 77568727



Period and Pulsation Analysis (DCDFT) for TIC 77568727



Light Curve for TIC 125754823



Period and Pulsation Analysis (DCDFT) for TIC 125754823



Light Curve for TIC 346626294



Period and Pulsation Analysis (DCDFT) for TIC 346626294



Light Curve for TIC 385552144



Period and Pulsation Analysis (DCDFT) for TIC 385552144



Light Curve for TIC 385553714



Period and Pulsation Analysis (DCDFT) for TIC 385553714



Light Curve for TIC 385558439



Period and Pulsation Analysis (DCDFT) for TIC 385558439



Light Curve for TIC 385589694



Period and Pulsation Analysis (DCDFT) for TIC 385589694



Light Curve for TIC 405483707



Period and Pulsation Analysis (DCDFT) for TIC 405483707



Light Curve for TIC 405483817



Period and Pulsation Analysis (DCDFT) for TIC 405483817



Light Curve for TIC 440690782



Period and Pulsation Analysis (DCDFT) for TIC 440690782



Light Curve for TIC 405484574



Period and Pulsation Analysis (DCDFT) for TIC 405484574



Light Curve for TIC 405461432



Period and Pulsation Analysis (DCDFT) for TIC 405461432



Light Curve for TIC 385509282



Period and Pulsation Analysis (DCDFT) for TIC 385509282



Light Curve for TIC 348639016



Period and Pulsation Analysis (DCDFT) for TIC 348639016



Light Curve for TIC 125754991



Period and Pulsation Analysis (DCDFT) for TIC 125754991



Light Curve for TIC 114083179



Period and Pulsation Analysis (DCDFT) for TIC 114083179



Light Curve for TIC 67768222



Period and Pulsation Analysis (DCDFT) for TIC 67768222



Light Curve for TIC 61139371



Period and Pulsation Analysis (DCDFT) for TIC 61139371



Light Curve for TIC 35155873



Period and Pulsation Analysis (DCDFT) for TIC 35155873



Light Curve for TIC 26126738



Period and Pulsation Analysis (DCDFT) for TIC 26126738



Light Curve for TIC 14111056



Period and Pulsation Analysis (DCDFT) for TIC 14111056

Discussion

The mechanism of the δ-Scuti pulsator is complex. Stellar pulsation modes directly probe a star's internal structure, including interior physical conditions and transport processes. The self-excited resonant pulsation modes of a star represent small perturbations to the hydrostatic equilibrium structure. Each pulsation is a standing wave with unique geometry, frequency, and pulsation signal structure including

harmonics, subharmonics, and signal higher order products that reflect nonlinear processes within the star. The frequencies can be separated into radial and angular components. Radial components are characterized by a radial order n, describing the number of interior shells acting as resonant cavities. Each resonant cavity supports a characteristic vibration mode. Angular components are characterized by an angular degree l, and an azimuthal order m, that describes the number of surface nodal lines. There are two main pulsation excitation mechanisms. The first is a stochastic mechanism that operates within large convective envelopes. Turbulent convection continually drives and damps the resonant frequencies of the star. These produce radial order pressure modes within certain frequency ranges determined by the resonant cavity. The second main pulsation excitement mechanism is a heat-engine mechanism in which mechanical work is converted to heat. This heats the various zones and causes them to expand. After expanding, the radiation is able to flow through the zones which releases the heat so the layers cool down and contract again. This periodic expansion-contraction cycle allows heat from the star to be converted into mechanical work and excite pulsations. Additionally, if the star is rapidly spinning, gravity can be important as a restoring force, resulting in additional frequencies and pulsation modes.

Conclusions

The Transiting Exoplanet Survey Satellite (TESS) observed pulsations in 34 eclipsing binary systems containing δ-Scuti stars in the instability strip of the Pleiades open cluster. Using time-series photometry data from TESS contained in the Mikulski Archive for Space Telescopes (MAST) database, light curves and periodograms are examined in detail to extract frequency/period relations among the pulsations, as well as the orbital periods of the binary components. In a few cases, the orbital period derived from the periodogram did not agree with those presented in Bedding, T. R., et al., 2023. No further analysis was pursued.

References

Bedding, T. R., et al., 2023, TESS Observations of the Pleiades Cluster: a Nursery for Delta Scuti Stars, arXiv:2212.12087v5, 18 March 2023

Benn, D. 2012, Algorithms + Observations = VStar, Journal of the AAVSO (JAAVSO), v40, n2, pp.852-866

Ferraz-Mello, S. 1981, Astron. J., 86, 619

Hinzel, D.H., 2025, The Transiting Exoplanet Survey Satellite (TESS) Observations of Eclipsing Binary V477 Cygni, JAAVSO Volume 53, 2025

Scargle, J. D., 1982, Astron. J., 263, 835

Space Telescope Science Institute (STScI). 2024, “Barbara A. Milulski Archive for Space Telescopes” MAST. https://archive.stsci.edu