

Time Domain Astrophysics

Multiwavelength, multimessenger, and time domain/transient events are increasingly important in astrophysics. Events to be studied include the neutrino precursors of Local Group Galactic Core Collapse Supernovae (CC-SN), optically bright Gravitational Wave (GW) events, and Gamma Ray Bursts (GRBs)/GRB afterglow. These events span widely different time frames, from a few per century (Local Group supernovae) to one per day for gamma-ray bursts. It would be advantageous if an integrated system could be implemented to detect the rare events as well as the more frequently occurring events.

Introduction and Background

There are several types or categories of high energy targets that would greatly benefit from follow-up optical observations from both professional and amateur astronomers. In addition to the Galactic Supernova, detected with the initial neutrino bursts, other potential high energy targets include Gamma Ray Bursts, GRB afterglow, and kilonovae. Additionally, there are several types of Gravitational Wave events such as neutron star-neutron star (NS-NS) mergers which can result in the formation of kilonovae, neutron star-black hole (NS-BH) mergers, and black hole-black hole (BH-BH) mergers. Each of these high energy events is based upon its own specific physics and detection mechanisms, but all of them require rapid optical follow-up observations. A brief description of the physics and detection mechanism for each type of high energy event follows.

Gamma Ray Bursts and Afterglow: In gamma-ray astronomy, gamma-ray bursts are extremely energetic explosions that have been observed in distant galaxies. They are the brightest electromagnetic events known to occur in the universe. Bursts can last from ten milliseconds to several hours. After an initial flash of gamma rays, a longer-lived "afterglow" is usually emitted at longer wavelengths (X-ray, ultraviolet, optical, infrared, microwave, and radio). The intense radiation of most observed GRBs is thought to be released during a supernova as a high-mass star implodes to form a neutron star or a black hole.

What are Gamma Ray Bursts (and why should we care?)

- Gamma radiation: Electro-magnetic radiation at wavelength even shorter than X-rays (boundary to X-rays is set arbitrarily)
- Blocked very efficiently by Earth's atmosphere
- Astrophysical gamma ray bursts first detected by military surveillance satellites (Vela)





Image Credits: NASA, Los Alamos National Laboratory

The sources of most GRBs are billions of light years away from Earth, implying that the explosions are both extremely energetic (a typical burst releases as much energy in a few seconds as the Sun will in its entire 10-billion-year lifetime) and extremely rare (a few per galaxy per million years). All observed GRBs have originated from outside the Milky Way galaxy, although a related class of phenomena, soft gamma repeater flares, is associated with magnetars within the Milky Way.

There are two kinds of gamma-ray bursts, known as long-soft and short-hard, referring to their duration and the nature of their gamma-ray emission. Long-soft bursts last for a few dozens of seconds, and emit less energetic ("soft") gamma rays; short-hard bursts last for a second or less and emit very energetic ("hard") gamma rays.

Two Flavors: Long & Short GRBs



Graph of burst duration versus the number of bursts. Graph of the time versus number of bursts for the gamma-ray bursts observed by the BATSE instrument on the Compton Gamma-ray Telescope.

Current understanding:

- Long duration GRBs: SN of very heavy stars
- Short duration GRBs: Neutron star mergers? ==> see later in this talk
- Connection to jets

Image Credit: NASA

The long-soft GRBs are the ones which have been detected most often at other wavelengths, and they are believed to be associated with the collapse of supermassive stars, in an event known as a hypernova. When a massive star runs out of the nuclear fuel that makes it shine, the core of the star collapses. If the core collapses into a black hole, the remainder of the star will begin to fall onto it. Black holes sometimes produce jets of material that fly away from the black hole at close to the speed of light, and in a hypernova, the infalling stellar material acts as a source for these jets. These events probably happen dozens of times a day across the entire universe, but we only detect them as a gamma-ray burst if, by chance, the jet from the black hole happens to be pointed in our direction. GRBs produce the most intense radiation along the direction of the jet, and so we only detect them when they're pointed right at us.


Although they haven't been studied as well, the short-hard GRBs are also believed to originate from the formation of a black hole. In this case it is thought that they come from the merger of two black holes or two neutron stars in orbit around one another. Both black holes and neutron stars are very massive and extremely small in size, and when they orbit one another closely, they move very rapidly. If they spiral together and merge with one another, their collision may result in a huge explosion that occurs very quickly, producing a rapid burst of gamma-rays at high energies.

Most of the energy emitted by a gamma-ray burst comes out as gamma-rays, but the jets that create them and the resulting hypernova emits light at other wavelengths

too, and by studying the afterglow, more can be learned about the object that created the GRB than can be from just studying the gamma-ray emission. The light emitted in X-rays, optical light, and radio waves can often persist for hours or days after the gamma-ray burst, and because of the nature of radiation at these wavelengths, it is easier to pinpoint where the GRB is from the afterglow than it is from the gamma-ray burst itself. It can be determined what kind of star it was that exploded, how the explosion progressed, or what the environment was like around that star by studying the afterglow.

GRB afterglows are hard to find, but there is now a network of space-based and ground-based observatories dedicated to their detection and localization. Satellites like Swift are designed to quickly detect and localize GRBs to much higher precision than was previously possible. Satellites can now provide gamma-ray localizations to less than 0.5 degrees (sometimes much less), making it easier for ground-based observers to concentrate their search on a particular spot in the sky. The satellite radios the coordinates back to Earth, and these coordinates are then relayed to observatories around the world via the Gamma Ray Burst Coordinates Network or GCN. Reference: [Gamma Ray Burst](#).


GRB detectors in space: FERMI (ex-GLAST)



- Two gamma ray detecting instruments
- **LAT**: FOV ca 20% of sky at any time, for all-sky survey
- **GBM**: together cover the entire sky not eclipsed by Earth
- Sky localization from GBM alone is usually poor (many arcminutes to degrees)
- Autonomous ToO slewing to better observe afterglow
- Onboard data analysis and alert generation

Image Credit: NASA

GRB detectors in space: Neil Gehrels Swift



- **Burst Alert Telescope:** Wide FOV (2 steradians), gamma ray low resolution “imager”
- Within 20 seconds of a GRB: 3 arcmin position estimate computed and sent to ground
- Then: Automatic slew of satellite to bring target into view of **X-Ray-Telescope** and **UV/Optical Telescope** to observe Afterglow
- Afterglow can then yield arcsec position and sometimes redshift

Image Credit: NASA

GRB detectors in space: Integral

INTERNATIONAL Gamma-Ray Astrophysics Laboratory (ESA)

- Main instrument (SPI) alone is not well suited for GRB detection
- SPI has a “stray gamma ray” detector (SPI-ACS) that can be used to detect GRBs





Image Credit: ESA

GRB detectors in space: The IPN

- Many deep space probes now carry simple and inexpensive gamma ray detectors
- Together they form the **InterPlanetary Network**
- Main contributing satellites: Konus-WIND*, 2001 Mars Odyssey, INTEGRAL, Swift, Fermi, and BepiColombo
- **Better localization** of GRBs (“triangulation”)...
- ...but much **higher latency**



GRB Afterglows (here's where amateurs come in)

- Afterglow: any emission following the GRB burst itself
- Also lower energy emission (X-rays, **optical**, radio)
- Interaction of relativistic jet with ISM? Synchrotron emission
- Not sure whether all GRBs have afterglows, maybe you just have to look fast and deep enough!

GRB021211 – optical detection by RAPTOR




image starting at:
 $t - t_0 = 64.9$ seconds

$t - t_0 = 9$ minutes

Credit: Image taken by the RAPTOR telescope and RAPTOR team at Los Alamos National Laboratory. Copyright 2002 LANL and the University of California

Gravitational Wave Events: Gravitational Wave events consist of three fundamental types of phenomena. These include Neutron Star-Neutron Star (NS-NS) mergers (also known as Binary Neutron Star or BNS mergers), Neutron Star-Black Hole (NS-BH) mergers, and Black Hole-Black Hole (BH-BH) mergers (also known as Binary Black Hole or BBH mergers). Additionally, kilonovae are associated with NS-NS GW events. A kilonova is a transient phenomenon that is relatively fast (days to weeks

timescale) and has a faint visible and infrared optical signature (this might also create a GRB). The radiation is fairly isotropic ejecta emitted at fractions of the speed of light (approximately 30%). The kilonova emissions are powered by the radioactive decay of heavy nuclei within the ejecta from intense bombardment of nuclei lighter than iron by energetic neutrons. Called the r-process, physics requires an energetic and extremely neutron-rich environment to be effective. The violent matter ejections resulting from the coalescence of two neutron stars can produce such an environment and have been proposed as being the main candidates of kilonovae and the production sites of the heaviest elements in the universe. Reference: [Gravitational Waves](#).

Gravitational Wave Detectors

Image credit: Caltech/MIT/LIGO Lab



LIGO Livingston,
Louisiana, USA

Virgo, Italy



Image credit: The Virgo collaboration/CCO 1.0


LIGO Hanford
Wash., USA



Image credit: Caltech/MIT/LIGO Lab

LIGO/Virgo Gravitational Wave Detectors

The Laser Interferometer Gravitational-Wave Observatory (LIGO) is a large-scale physics experiment and observatory to detect cosmic gravitational waves and to develop gravitational wave observations as an astronomical tool. Two large observatories were built in the United States (Hanford, Washington and Livingston, Louisiana) with the aim of detecting gravitational waves by laser interferometry. These observatories use mirrors spaced four kilometers apart which are capable of detecting a change of less than one ten-thousandth the charge diameter of a proton. LIGO detectors use laser interferometry to measure the distortions in space-time occurring between stationary, hanging masses (mirrors) caused by passing gravitational waves. The distortions in space-time were predicted by Einstein's General Theory of Relativity. This is shown below. Reference: [LIGO](#).



LIGO Gravitational Wave Detector (Abbot, B.P., et.al, LIGO Scientific Collaboration and Virgo Collaboration)

Advanced Virgo is an interferometric detector of gravitational waves hosted by the European Gravitational Observatory (EGO) near Pisa, Italy. It can measure gravitational waves in a wide frequency range; from 10 Hz to 10000 Hz. Advanced Virgo is a laser interferometer with perpendicular, 3 km long arms and suspended mirrors. This is the configuration presently adopted in Advanced Virgo, during the third observation run (named O3, lasting 1 year from April 2019 to April 2020) performed jointly with the two Advanced LIGO detectors in the US. The Virgo system is similar to that of LIGO. Reference: [VIRGO](#).


Gravitational Wave detection events are transmitted via GCN as discussed below for gamma ray bursts/afterglows. The first gravitational wave event detection and localization included a gravitational wave (GW 170817A), a GRB (GRB 170817), and a subsequent kilonova (AT2017gfo).

First Detection: GW 170817A GRB170817 & AT2017gfo

- Aug 17 2017 :
A Gravitational wave event is recorded by LIGO
- At the same time (~1.5 sec later), a GRB is detected by Fermi and Integral

Image credit: Gravitational Waves and Gamma-Rays from a Binary Neutron Star Merger: GW170817 and GRB 170817A, Fig. 2
<http://iopscience.iop.org/article/10.3847/2041-8213/aa920c>
 DOI: 10.3847/2041-8213/aa920c
 License Creative Commons Attribution 3.0 Unported

02/08/2022



First Detection: GW 170817A GRB170817 & AT2017gfo

- The GW detectors and Fermi provide an approximate sky-localization
- Less than 11h after the event, an optical transient is found: the kilonova AT2017gfo

THE ASTROPHYSICAL JOURNAL LETTERS, 848:L12 (9pp), 2017 October 20

Abbott et al.





Figure 1. Localization of the gravitational-wave, gamma-ray, and optical signals. The left panel shows an orthographic projection of the 90% credible regions for LIGO (190 deg²; light green), the initial LIGO-Virgo localization (131 deg²; dark green), IPN triangulation from the time delay between Fermi and INTEGRAL (light blue), and Fermi-GBM (dark blue). The inset shows the location of the apparent host galaxy NGC 4993 in the Swope optical discovery image at 10.9 hr after the merger (top right) and the DLT40 pre-discovery image from 28.5 days prior to merger (bottom right). The reticle marks the position of the transient in both images.

Original content from this work may be used under the terms of the Creative Commons Attribution 3.0 license. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

Galactic Core Collapse Supernova: These are extremely rare events, occurring 1-10 times per century. As such, it is important to have the technical capability to detect galactic supernovae and quickly perform as much follow-on optical backup as possible. When a massive star reaches the end of its life, its core collapses. More than 99% of the binding energy of the resulting neutron star or black hole is released in the form of neutrinos and antineutrinos of all flavors, with energies in the tens of MeV (million electron volt) range. The energy leaves via neutrinos/antineutrinos because


neutrinos/antineutrinos interact so weakly that they readily leave the star. They bring information from deep inside the stellar core and great insight about core collapse physics. The time scale of neutrino/antineutrino emission is a few tens of seconds immediately after core collapse. The optical photon signal, in contrast, can take hours to days to emerge from the stellar envelope. Therefore, the detection of a burst of neutrinos/antineutrinos will give an early warning of the supernova collapse so that optical followup will be possible. Reference: Scholberg, K., [\[0803.0531\] The SuperNova Early Warning System](#).



[Observing the Next Galactic Supernova](#)

Core collapse supernova detection does not use spacecraft as in the case of GRBs/afterglows. The basic idea of the core collapse supernova alert system is to have a central computer which accepts neutrino burst candidate messages from neutrino detectors around the world and sends an alarm message to astronomers if it finds a coincidence within a short period of time (10 seconds). The central computer is located at Brookhaven National Lab. The coincidence search is both “blind” (decision is made when messages are received without polling the other experiments) and automated (alerts go out without human intervention for maximum speed). The neutrino experiments currently involved are Super-K, LVD, IceCube, Borexino, KamLAND, Daya Bay, and HALO. This is depicted below along with the interface to GCN for followup electromagnetic/optical observations. Reference: [SNEWS](#) and <https://snews2.org/>.

SNEWS System Architecture




High Energy Event Detection Triggers and Alerts

GRBs/afterglow, Gravitational Wave events, and Core Collapse Supernovae all utilize the GRB Coordinate Network (GCN) hosted by the NASA Goddard Space Flight Center (GSFC).

GCN architecture

The GRB observational technique begins with the distribution of information from the [Gamma-ray Coordinate Network](#), specifically the [GCN Circular Archive](#). There are two parts to the GRB Coordinates Network: (1) the distribution of GRB/transient locations detected by various spacecraft, and (2) receiving and automatically distributing to the GRB/transient community prose-style e-mail messages about follow-up observations on various GRBs/transients. The overall high level physical GCN architecture is shown below.



The Physical GCN Network (Courtesy NASA Goddard Space Flight Center)

GCN Circular Archive

Part 1 (the GCN Notices):

This portion of GCN consists of distributing the GRB/transient locations determined by the Fermi, Swift, INTEGRAL, IPN (Interplanetary Network), MOA (micro-lensing events), and other spacecraft systems. The collection and distribution of these Notices are done without any humans-in-the-loop, and as such for missions with real-time downlinks, the time delay from when the GRB/transient occurs to when the Notices are being sent out to the customer/observer is in the 2-10 sec range. Many socket-based and email-based formats and protocols are available. This combining of all the sources of GRB/transient location information into a single network means that user sites need only maintain a single interface for all their GRB/transient needs.

Example GCN notice (in human readable form)

```
//////////////////////////////////////
TITLE:          GCN/SWIFT NOTICE
NOTICE DATE:    Tue 18 Jan 22 18:30:12 UT
NOTICE TYPE:    Swift-UVOT Position
TRIGGER_NUM:    1093717, Seg Num: 0
GRB_RA:         192.2688d (+12h 49m 04.51s) (J2000),
                192.5482d (+12h 50m 09.64s) (current),
                191.6528d (+12h 40m 36.67s) (1950)
GRB_DEC:        +22.9156d (+22d 54' 56.1") (J2000),
                +22.7957d (+22d 47' 44.5") (current),
                +22.1000d (+23d 11' 16.6") (1950)
GRB_EPOCH:      0.8 [arcsec radius, statistical only]
GRB_MAG:        17.82 +/- 0.14 [mag]
FILTER:         10.0micron
IMG_START DATE: 19597 TJD; 18 DOY; 22/01/18
IMG_START TIME: 66134.00 SOD (18:22:14.00) UT, 95.9 [sec] since BAT Trigger Time
SUN_POSTN:      300.77d (+20h 03m 04s) -20.43d (-20d 25' 42")
SUN_DIST:       113.92 [deg] Sun_angle= 7.2 [hr] (West of Sun)
MOON_POSTN:     131.14d (+08h 44m 33s) +22.90d (+22d 54' 08")
MOON_DIST:      56.13 [deg]
MOON_ILLUM:     99 [%]
GAL_COORDS:     295.57, 85.75 [deg] galactic lon,lat of the burst (or transient)
ECL_COORDS:     181.57, 25.79 [deg] ecliptic lon,lat of the burst (or transient)
COMMENTS:       SWIFT UVOT Position Notice.
COMMENTS:       This Notice was ground-generated -- not flight-generated.
COMMENTS:       The UVOT position is 10.8 arcsec from the XRT position.
COMMENTS:       Result based on Genie data.
COMMENTS:       Notice generated automatically.
```



Allows **automatic**
decision making:

*Is this event observable
from my site with my
equipment this night*

?

Or:

*Automate your
robotic telescope!*

Courtesy NASA Goddard Space Flight Center

Part 2 (the GCN Circulares):

This part allows the GRB community to submit messages to a central queue where they are automatically vetted and distributed (via email) to the entire GRB community. These are prose-style messages (as opposed to the "TOKEN: value" style of the GCN e-mail Notices) from follow-up observers reporting on their results (detections or upper limits) or for coordinating with others. Reference: [GCN](#).

Public GCN circulars on the Internet

GCN circulars can be subscribed to by email, and you can see them here:

https://gcn.gsfc.nasa.gov/gcn3_archive.html

GCN Circulars Archive (in serial number order)

This page changes after each Circular submission, so hit the <reload> button NOW.

The processing of Circulars to this page was stopped (from ~01:30 to ~18:00 UT 01 Dec 2020).
The Circulars page has been fixed as of ~18:00 UT 01 Dec 2020.
(The Circulars WERE ALWAYS BEING DISTRIBUTED -- it is only this archive web page (and sub-pages) that had problems.)

1. The Latest Circulars
2. Older Circulars
3. Torfile of all Circulars
4. Circulars grouped by each Event
5. All Circulars on the GRB source type
6. All Circulars on the GW source type
7. All Circulars on the SGR source type
8. All Circulars on the misc source type (ie not in any of the above 3 pages).

This is the archive of all the GCN Observation Report Circulars (listed in reverse serial number order, newest first).
To learn more about the [GCN Circulars](#) and [how to subscribe](#) and to [unsubscribe here](#).

1) The Latest Circulars (listed in reverse serial number order -- newest first)

- 31541 GRB 220131B: Swift/BAT-GUANO detection outside the coded FOV
- 31540 GRB 220201A: Fermi GBM Final Localization
- 31539 GRB 220128A: AstroSat CZTI detection
- 31538 GRB 220130A: Fermi GBM Observation
- 31537 Konus-Wind detection of GRB 220128A
- 31536 IPN triangulation of GRB 220128A (long)
- 31535 Fermi GRB 220131A: Global MASTER-Net observations report
- 31534 GRB 220131B: GECAM detection

Example GCN circular

```
////////////////////////////////////  
TITLE:   GCN CIRCULAR  
NUMBER:  31490  
SUBJECT: GRB 220118A: Swift detection of a burst with an optical counterpart  
DATE:    22/01/18 18:33:38 GMT  
FROM:    Kim Page at U.of Leicester <kimlpage1978@gmail.com>
```

N. J. Klingler (GSFC/UMBC/CRESSTII), E. Ambrosi (INAF-IASFPA),
A. D'Ai (INAF-IASFPA), V. D'Elia (SSDC & INAF-OAR), S. Dichiara (PSU),
J.D. Gropp (PSU), H. A. Krimm (NSF), F. E. Marshall (NASA/GSFC),
K. L. Page (U Leicester), D. M. Palmer (LANL),
T. M. Parsotan (GSFC/UMBC/CRESSTII), B. Sbarufatti (PSU) and
M. H. Siegel (PSU) report on behalf of the Neil Gehrels Swift
Observatory Team:

At 18:20:38 UT, the Swift Burst Alert Telescope (BAT) triggered and
located GRB 220118A (trigger=1093742). Swift slewed immediately to the burst.
The BAT on-board calculated location is
RA, Dec 152.274, +22.909 which is
RA(J2000) = 12h 49m 06s
Dec(J2000) = +22d 54' 23"
with an uncertainty of 3 arcmin (radius, 90% containment, including
systematic uncertainty). The BAT light curve showed a single-peaked
structure with a duration of about 20 sec. The peak count rate
was ~1400 counts/sec (15-350 keV), 0.5 sec after the trigger.

13 minutes
delay

Courtesy NASA Goddard Space Flight Center

Example GCN circular cont'd

The XRT began observing the field at 18:22:08.3 UT, 90.3 seconds after the BAT trigger. Using promptly downlinked data we find a bright, fading, uncatalogued X-ray source located at RA, Dec 192.26936, 22.91000 which is equivalent to:

RA(J2000) = 12h 49m 04.65s
Dec(J2000) = +22d 54' 57.6"

with an uncertainty of 3.5 arcseconds (radius, 90% containment). This location is 29 arcseconds from the BAT onboard position, within the BAT error circle. This position may be improved as more data are received; the latest position is available at <https://www.swift.ac.uk/sper>.

A power-law fit to a spectrum formed from promptly downlinked event data gives a column density in excess of the Galactic value ($2.25 \times 10^{20} \text{ cm}^{-2}$, Willingale et al. 2013), with an excess column of $3.4 (+2.60/-2.24) \times 10^{21} \text{ cm}^{-2}$ (90% confidence).

The initial flux in the 2.5 s image was $4.57\text{e-}10 \text{ erg cm}^{-2} \text{ s}^{-1}$ (0.2-10 keV).

Courtesy NASA Goddard Space Flight Center

Example GCN circular cont'd

UVOT took a finding chart exposure of 150 seconds with the White filter starting 99 seconds after the BAT trigger. There is a candidate afterglow in the rapidly available $2.7' \times 2.7'$ sub-image at

RA(J2000) = 12:49:04.53 = 192.26888
DEC(J2000) = +22:54:56.2 = 22.91552

with a 90%-confidence error radius of about 0.75 arc sec. This position is 1.8 arc sec from the center of the XRT error circle. The estimated magnitude is 17.83 with a 1-sigma error of about 0.15. No correction has been made for the expected extinction corresponding to E(B-V) of 0.025.

Burst Advocate for this burst is N. J. Klingler (noelklin AT umbc.edu). Please contact the BA by email if you require additional information regarding Swift followup of this burst. In extremely urgent cases, after trying the Burst Advocate, you can contact the Swift PI by phone (see Swift T00 web site for information: <http://www.swift.psu.edu/>)

Courtesy NASA Goddard Space Flight Center

GRB Afterglow Amateur Observations

- **Filipp Romanov**
(obs code RFDA)

Example:



- Read about in in the AAVSO HEN Forum
- Images => flickr stream

<https://www.flickr.com/photos/filipp-romanov/>

TITLE: GCN CIRCULAR
 NUMBER: 26505
 SUBJECT: GRB 191221B: iTelescope optical observations
 DATE: 19/12/23 00:21:24 GMT
 FROM: Filipp Dmitrievich Romanov at Amateur astronomer <filipp.romanov@iTelescope.net>

I observed the optical afterglow of GRB 191221B (Laha et al., GCN Circ. 26534) with remote telescope T32 (0.43-m f/6.8 reflector + FLI Proline 16803 CCD) of iTelescope.Net in Siding Spring Observatory (Australia) on 2019-12-22 since 13:28:52 UT (~16.8 hours after the trigger). Two images were obtained in each band with Astrodon luminance filter and Johnson/Cousins V filter for 180 and 120 seconds (Blnx2). I detected the optical afterglow in three images in the enhanced Swift-XRT position (Beardmore et al., GCN Circ. 26549); it not present in images from DSS Plate Finder.

The following magnitudes (mag. error ~ 0.2) were measured from comparison to nearby stars from APASS DR9 catalogue (Henden et al., 2016):

Time of start	Exp.time	Mag.	Comp.star	band	Filter
13:28:52 UT	180 s	18.1	r'		Luminance
13:32:58 UT	180 s	18.3	V		V
13:36:51 UT	120 s	18.3	r'		Luminance
13:39:46 UT	120 s	<17.9	V		V

Magnitudes were not corrected for Galactic extinction.

Courtesy Filipp Romanov

GRB Afterglow Amateur Observations

- **Arto Oksanen**
(obs code OAR)
- Discovered & co-discovered several GRB afterglows
- E.g. read about it in :



Oksanen et al., *AAVSO Volume 36, 2008* 53

Discovery and Observations of the Optical Afterglow of GRB 071010B

Arto Oksanen
Hankasalmi Observatory, Hankasalmi, Finland

Matthew Templeton
Arne A. Henden
AAVSO, 49 Bay State Road, Cambridge, MA 02138

David Alexander Kann
Thüringer Landessternwarte Tautenburg, Sternwarte 5, D-07778 Tautenburg, Germany

Received December 18, 2007; revised February 11, 2008; accepted February 11, 2008

Abstract On 2007 October 10 at 20:45:48 UT, the Swift satellite detected the bright, long-soft gamma-ray burst GRB 071010B in the constellation Ursa Major. Coordinates were automatically distributed via the Gamma-ray Burst Coordinate Network (GCN), and observations were begun by A. Oksanen at the Hankasalmi Observatory in Hankasalmi, Finland, within fifteen minutes of the burst. A previously uncatalogued optical source was detected at R.A. 10^h 02^m 09.26^s, Dec. +45° 43' 50.3" (J2000) at an unfiltered (R-band calibrated) magnitude of approximately 17.5. Imaging over the following six

02/08/2022

Courtesy Arto Oksanen

Time Domain Astrophysics Techniques

GCN is utilized for all three types of time domain events. In order to interface with the GCN, the observer simply needs anything that will run Python code and has a 24/7 internet connection. Because the messages are machine readable and the API is Python, the observer can react to events in a very customizable way, e.g., you can filter out “interesting” events based on observability at your site or the size of the uncertainty of the sky localization. All it takes to receive these messages are a few lines of Python code, thanks to the PyGCN package that you can find here: <https://pypi.org/project/pygcn/>. There are two servers available (identical servers for redundancy); 45.58.43.186 port=8099 and 68.169.57.253 port=8099. These addresses are already preconfigured in the Python library as defaults <https://github.com/lpsinger/pygcn>.

PyGCN example: listening for FERMI:

```
#!/usr/bin/env python
import gcn

# Define your custom handler here.
@gcn.include_notice_types(
    gcn.notice_types.FERMI_GBM_FLT_POS, # Fermi GBM localization (flight)
    gcn.notice_types.FERMI_GBM_GND_POS, # Fermi GBM localization (ground)
    gcn.notice_types.FERMI_GBM_FIN_POS) # Fermi GBM localization (final)
def handler(payload, root):
    # Look up right ascension, declination, and error radius fields.
    pos2d = root.find('://{*}Position2D')
    ra = float(pos2d.find('://{*}C1').text)
    dec = float(pos2d.find('://{*}C2').text)
    radius = float(pos2d.find('://{*}Error2Radius').text)

    # Print.
    print('ra = {:g}, dec={:g}, radius={:g}'.format(ra, dec, radius))


# Listen for VOEvents until killed with Control-C.
gcn.listen(handler=handler)
```

PyGCN example: listening for SNEWS:

```
#!/usr/bin/env python
import gc

# Define your custom handler here.
@gcn.include_notice_types(
    gc.notice_types.SNEWS)
def handler(payload, root):
    print('SNEWS Alert!')
    # .... if at night: switch bedroom lights on
    # .... play a loud sound
    # .... send emails to yourself and friends
    # P.S. there are test alerts every Tuesday , you might want
    # to filter out those once you are confident everything
    # works ... :-)
# Listen for VOEvents until killed with Control-C.
gc.listen(handler=handler)
```

Raspberry Pi: Raspberry Pi is an inexpensive, single board microcomputer that can be programmed for a wide variety of applications. One such application is an alert system that receives, processes, and displays GCN alerts from LIGO/VIRGO, Goddard Space Flight Center, and any detector/trigger that requires a rapid follow-up response (e.g., an optical observation of a Gamma Ray Burst or Gravitational Wave Event). Raspberry Pi can be utilized by amateur observers and/or robotic telescope systems to configure their equipment to perform observations. PyGCN, as described above, can be run on Raspberry Pi.



Courtesy Heinz-Bernd Eggenstein

Mobile/Cell Phone Apps (iPhone/Android): One potentially important high energy event alert is mobile apps for cell phones, both iPhone and Android. These apps receive a GCN or similar high energy event trigger and generate an SMS/text message to a user's mobile device. This would likely be one of the quickest ways to alert observers that an event has occurred, since cell phone use is ubiquitous. There are two unofficial apps for mobile devices that can be used to view GW alerts on smartphones or tablets. These can be found here [Mobile Device GW Alerts](#).

GRB Observational Techniques

1. Receipt of GCN message: GCN sends a message, as described above, from the Swift gamma-ray location function. Coordinates are automatically distributed in the message. The detected GRB/afterglow should be observed as soon as possible after receiving the GCN alert. It is also useful to learn to know if/when a GRB has potential prior to any alert.
2. Telescope observations: In principle, GRBs can be imaged within seconds of detection of gamma-rays so long as the observer is in darkness, can view the field from their location, has a clear sky, and has equipment quickly available (a large telescope with CCD camera helps). The telescope characteristics include the location (latitude/longitude/elevation) of the observing platform as well as the

optical and photometric parameters of the telescope. For example, iTelescope T18, AstroCamp Observatory, Nerpio, Spain (North 38 deg/ West 2 deg, 19 min/ 1650 meters); 0.32 m, f/8.0 reflector + SBIG-STXL-6303E CCD. Some specific telescope systems used to develop these observation procedures are:

[iTelescope](#)

iTelescope is a worldwide network of robotically controlled telescopes. It is comprised of four observatories in both the Northern and Southern Hemispheres: Siding Springs Observatory, Australia; Astrocamp, Nerpio, Spain; New Mexico Skies, New Mexico; and Sierra Remote Observatory, California. These observatories host 22 robotic telescopes with varying capabilities controlled across the internet.

[Abbey Ridge Observatory](#)

The observatory is built around a fiberglass 10-foot diameter [Home-Dome](#). Inside is a [Celestron](#) C14 Schmidt-Cassegrain Telescope mounted on a Losmandy [Titan](#) German equatorial mount controlled by a SiTech controller. Its imaging camera is an [SBIG](#) ST8XME CCD camera with dual Optec IFW filter wheels (9 effective filters), Optec NextGen telecompressor, and an Optic TCF electric focuser. Abbey Ridge Observatory also uses an AAG CloudWatcher to monitor the sky conditions. There are more details [here](#).

[Burke-Gaffney Observatory](#)

The Burke-Gaffney Observatory and its [Dr. Ralph M. Medjuck](#) Telescope are primarily used for undergraduate astronomy education, public outreach, and modest research projects. It is one of only two observatories in the world, along with the nearby [Abbey Ridge Observatory](#), that can be [controlled from Twitter, Facebook Messenger, cell phone text messaging, and a dedicated App](#) in a fully-automatic way.

The observatory is named in honor of [Reverend M. W. Burke-Gaffney](#) , S. J (1896-1979). The new telescope, installed in December of 2013, is named in honor of [Dr. Ralph M. Medjuck](#). It is located on the roof of the 22-story Loyola Residence tower. It was opened in 1972 with a 16-inch Ealing telescope, but was extensively upgraded and renovated in 2013. The main instrument is a Planewave CDK24, a 0.61-metre diameter reflecting telescope. Technical details for the telescope and its instruments can be found [here](#). The observatory also owns a 40-cm diameter reflecting telescope, ten 20-cm diameter reflecting telescopes (with several white light solar filters), and a dedicated solar telescope with a narrowband hydrogen alpha filter.

3. Observation and Image Parameters: The observation parameters include UT time of exposure, time after trigger (for example 2020-12-23 from 19:49:22 UT: 1 hour, 50 minutes, 56 seconds after trigger), RA/Dec of observations, e.g., RA

08:51:09.46, Dec +71:10:47:00. Image parameters include number of images, exposure time, bin number, and filter type (e.g., V, luminance, Sloan). For example, 5 images with exposures 300, 120, 180 seconds with an Astrodon luminance filter and 4 images with exposures 300 and 60 seconds with a photometric Johnson V filter. Finally, any image stacking should be indicated, for example, 4x300 + 2x120 + 180 seconds. More specifically, 2x300V + 2x300lum + 2x120lum + 180lum seconds.

4. Results: Indicate all of the pertinent results of the observations and measurements. For example, measured magnitude from comparison to r' magnitudes of nearby stars from the Pan-STARRS DR1 catalog (Chambers et al., 2016). This will typically be as follows:


Time of start 19:49:22 UT; Exposure time 300 seconds; Bin=1; Mag. r' =19.2; Mag error 0.2; Filter = Luminance

Time of start 19:56:25 UT; Exposure time 300 seconds; Bin=1; Mag. r' =19.3; Mag error 0.4; Filter = V

Time of start 20:26:25 UT; Exposure time 300 seconds; Bin=2; Mag. r' =19.9; Mag error 0.4; Filter = V

Time of start 20:33:07 UT; Exposure time 300 seconds; Bin=2; Mag. r' =20.1; Mag error 0.3; Filter = Luminance

Include magnitude, error, comparison star band, limits of parameters if applicable, and SNR. Indicate if a correction was made for galactic extinction (yes/no) and indicate if an optical afterglow was detected (yes/no). Finally, present and publish the finished optical image product and supporting data on the [AAVSO International High Energy Network HEN Forum](#). An example of a finished image product is:



Courtesy Filipp Romanov

Include any appropriate notes, information, data, comments, suggestions for further observations or research, etc.

5. Submit a GCN Circular of the discovery/observation as soon as possible with all the pertinent data. Here are links to sample GCN Circulars (courtesy of Filipp Romanov and Arto Oksanen). If assistance is needed, contact Arto Oksanen at arto.oksanen@jkl Sirius.fi.

Sample GRB Circulars (Courtesy Filipp Romanov and Arto Oksanen)

GRB 191029A: iTelescope optical upper limits

<https://gcn.gsfc.nasa.gov/gcn/gcn3/26091.gcn3>

GRB 191123A: iTelescope optical upper limits

<https://gcn.gsfc.nasa.gov/gcn/gcn3/26283.gcn3>

GRB 191221B: iTelescope optical observations

<https://gcn.gsfc.nasa.gov/gcn/gcn3/26565.gcn3>

GRB 201223A: iTelescope optical afterglow observations

<https://gcn.gsfc.nasa.gov/gcn/gcn3/29165.gcn3>

GRB 210104A: iTelescope optical upper limit

<https://gcn.gsfc.nasa.gov/gcn/gcn3/29269.gcn3>

GRB 210226A: Burke-Gaffney Observatory optical upper limit

<https://gcn.gsfc.nasa.gov/gcn/gcn3/29572.gcn3>

GRB 210306A: iTelescope optical afterglow observations

<https://gcn.gsfc.nasa.gov/gcn/gcn3/29599.gcn3>

GRB 210610B: iTelescope optical afterglow observations

<https://gcn.gsfc.nasa.gov/gcn/gcn3/30181.gcn3>

GRB 210610B: correction to GCN 30181

<https://gcn.gsfc.nasa.gov/gcn/gcn3/30184.gcn3>

GRB 210610B: optical observations from Burke-Gaffney Observatory and Abbey Ridge Observatory

<https://gcn.gsfc.nasa.gov/gcn/gcn3/30228.gcn3>

GRB 210619B: iTelescope optical afterglow observation

<https://gcn.gsfc.nasa.gov/gcn/gcn3/30292.gcn3>

GRB 210619B: Abbey Ridge Observatory optical afterglow observation
<https://gcn.gsfc.nasa.gov/gcn3/30305.gcn3>

GRB 210702A: iTelescope optical afterglow observations
<https://gcn.gsfc.nasa.gov/gcn3/30364.gcn3>

GRB 220101A: iTelescope optical observations
<https://gcn.gsfc.nasa.gov/gcn3/31387.gcn3>

GRB 210822A: Abbey Ridge Observatory optical afterglow observation
<https://gcn.gsfc.nasa.gov/gcn3/30701.gcn3>

GRB 210702A: iTelescope optical afterglow observations
<https://gcn.gsfc.nasa.gov/gcn3/30364.gcn>

GRB 071010B possible optical transient
[GCN Circular 6873](#)

GRB140206A - Optical afterglow candidate
[GCN Circular 15786](#)

GRB 071010B correction to GCN 6892 optical observations
[GCN Circular 6903](#)

Gravitational Wave Observational Techniques

The interfaces for the amateur optical follow-up of gravitational wave events are GRANDMA and Kilonova Catcher.

[Global Rapid Advanced Network Devoted to the Multi-messenger Addicts \(GRANDMA\)](#)

An international scientific collaboration of astrophysicists that work on gravitational wave astronomy, utilizing crowdsourcing and volunteer astronomers to assist in research. GRANDMA utilizes a network of robotically controlled telescopes with both photometry and spectrometry capabilities for time-domain astronomy.


Kilonova Catcher

Kilonova Catcher is focused on the optical followup of any gravitational wave event that is detected by the currently operating gravitational wave observatories, specifically to catch the [kilonova](#) emissions emerging from the coalescence of two neutron stars bound in a compact binary system.

Kilonova Catcher Project

- <https://grandma-kilonovacatcher.lal.in2p3.fr/>


- Pro-Am-Collaboration
- Watch the AAVSO Webinar, 2nd part of <https://youtu.be/WIDxZDA2ZvY>



Watch the AAVSO Webinar, 2nd part, <https://youtu.be/WIDxZDA2ZvY> for a detailed discussion of GRANDMA/Kilonova Catcher with Dr. Sarah Antier.

The overall system architecture of GRANDMA/Kilonova Catcher is shown below.

GRANDMA-Kilonova Catcher




Courtesy Kilonova Catcher

Galactic Supernova Observational Techniques

The SuperNova Early Warning System (SNEWS) is the interface for the amateur optical follow-up of Core Collapse Supernovae. SNEWS sends neutrino detection alerts in real time via email or NASA's GCN system. The initial alert is currently without sky localization.

SNEWS 2.0 : visit <https://snews2.org>



Reference: [SNEWS](https://snews2.org/) and <https://snews2.org/> and <https://snews2.org/alert-signup/>

Watch the AAVSO Webinar, 3rd part, <https://youtu.be/WIDxZDA2ZvY> for a detailed discussion of Gravitational Waves and Multi-messenger astronomy with Dr. Michael Coughlin.


How amateurs can and should prepare for observing these events

Different ways to contribute, ALL useful!

- Discovery



- "Precovery"



- Follow-up

One way to monitor possible time domain events is through the Astronomer's Telegram (<https://www.astronomerstelegram.org/>).

The Astronomer's Telegram
 5 Feb 2022; 15:48 UT

Telegram Index
 Telegrams Posted Within the Last 30 Days (All)
 50 Selected of 15201 Telegrams

15201 No bright optical counterpart to the GeV transient; Fermi J0614+1713

15200 Observations of the new gamma-ray transient Fermi J0614+1713 with VERITAS

15199 Improved Localization of the Fermi-LAT Gamma-ray Transient in the Galactic Plane; Fermi J0614+1713

15198 H-alpha observations of M31 nova candidates AT2022air and AT2022iuz

15197 Detection of two bright bursts from FRB20201124A with AperiT of the Westerbork Synthesis Radio Telescope.

15196 Fermi LAT Detection of a New Gamma-ray Transient in the Galactic Plane; Fermi J0614+1713

Recently
15194 Optical follow-up of MAXI J0709-159 (B08010101)
15193 NuSTAR identified MAXI J0709-159 as a No. X-ray binary with LX ON/OFF
15188 MAXI observations of two bright X-ray flares from MAXI J0709-159
15181 NuSTAR detection of the new X-ray transient MAXI J0709-159
15178 HAWKINE detection of an X-ray short transient associated with MAXI J0709-159

Most Viewed
15200 Observations of the new gamma-ray transient Fermi J0614+1713 with VERITAS
15199 Improved Localization of the Fermi-LAT Gamma-ray Transient in the Galactic Plane; Fermi J0614+1713
15201 No bright optical counterpart to the GeV transient Fermi J0614+1713

Additionally, observers can access the Astro Colibri App Alert and Observation Tool.

<https://astro-colibri.herokuapp.com/welcome>


<https://astro-colibri.com/>

Astro Colibri App: Alert and Observation Tool



IOS and Android apps in their respective app stores can send notifications to your phone for new events including GRBs, GW events, as well as supernova and other transients.

Additionally, by monitoring red supergiant stars ([Red Supergiants](#)), it may be possible to build the light curve history in order to understand if a star might be a candidate for explosion. The AAVSO will manage the candidate list and make it available as a web page and an API to facilitate the effort.



Concept and implementation by
Lauren Herrington

Example Walk-Through

GCN

Example Walk-Through

- GRB 220124A

TITLE: GCN CIRCULAR
NUMBER: 31521
SUBJECT: GRB 220124A: AstroSat CZTI detection of a bright long GRB
DATE: 22/01/24 08:11:18 GMT
FROM: Gaurav Waratkar at IIT,Bombay <gauravwaratkar@iitb.ac.in>

V. Prasad (IUCAA), G. Waratkar (IITB), A. Suresh (IITB), A. Vibhute (IUCAA), V. Bhalerao (IITB), D. Bhattacharya (IUCAA), A. R. Rao (IUCAA/TIFR), and S. Vadawale (PRL) report on behalf of the AstroSat CZTI collaboration:

Analysis of AstroSat CZTI data with the ML pipeline (Abraham et al., 2021, MNRAS, 504, 3084) and the CIFT framework (Sharma et al., 2021, JApA, 42, 73) showed detection of a long GRB 220124A.

Example Walk-Thru

- However....

We note that the burst has not been localised, and we cannot rule out an association with any SGRs. The ASIMOV tool (<http://astrosat-ssc.iucaa.in:8080/ASIMOV/ASIMOV> [1]) indicates that other active missions were covering different parts of the sky, hence confirmed non-detections from other missions can help localise the source by inferring earth-occulted regions.

Example Walk-Thru

- Later on GCN:

GECAM also detected the GRB, and it has a localization

```
TITLE: GCN CIRCULAR
NUMBER: 31524
SUBJECT: GRB 220124A: GECAM detection
DATE: 22/01/25 14:48:59 GMT
FROM: Y Q Zhang at IHEP <yqzhang@ihep.ac.cn>
```

```
J. C. Liu, S. L. Xiong, S. Xiao, Y. Q. Zhang, C. Cai, P. Zhang,
C. Y. Li, S. L. Xie, X. Y. Zhao, Y. Huang, X. Y. Song,
C. Zheng, Y. Zhao, Z. W. Guo, W. C. Xue, C. W. Wang,
Q. B. Yi, B. X. Zhang, W. X. Peng, R. Qiao, D. Y. Guo, X. B. Li,
X. Ma, L. M. Song, P. Wang, J. Wang, Z. Zhang, S. J. Zheng, W. Chen,
J. J. He, G. Y. Zhao, Y. Q. Du, H. Wu, J. Liang, Q. Luo, X. L. Zhang,
H. M. Zhang, Z. H. An, M. Gao, K. Gong, B. Li, C. Li, J. H. Li,
X. Q. Li, Y. G. Li, X. H. Liang, X. J. Liu, Y. Q. Liu, X. L. Sun,
Y. L. Tuo, J. Z. Wang, X. Y. Wen, Y. B. Xu, Y. P. Xu, S. Yang,
C. Y. Zhang, D. L. Zhang, Fan Zhang, Fei Zhang,
X. Zhou, F. J. Lu, S. N. Zhang (IHEP)
report on behalf of GECAM team:
```

During the commissioning phase, GECAM-B was triggered in-flight by a long burst, GRB 220124A, at 2022-01-24T03:37:38.200 UTC (denoted as T0), which was also observed by CALET/GBM(triggerID: 1327030648), AstroSAT/CZTI(GCN #31521).

GECAM alert data was downlinked to the ground through the short message service of BeiDou Navigation Satellite System (BDS) within ~60 s after T0.

According to the BDS alert data, this burst mainly consists of multiple pulses with a duration of about 240 s.

The GECAM light curve could be found here:
<http://ftp.astro.ihep.ac.cn/pub/astromin/GRB220124ALC.png>

Using the automatic on-ground localization pipeline with the BDS alert data, GECAM-B localized this burst to the following position (J2000):

Ra: 246.18 deg
Dec: 42.29 deg


Err: 2.96 deg (1-sigma, statistical only)

The current systematic error of location is estimated to be several degrees and could be minimized by the ongoing observation.

GCN

Example Walk-Thru

- Still later on GCN: IPN has a “triangulation out on GCN:



TITLE: GCN CIRCULAR
NUMBER: 31525
SUBJECT: IPN triangulation of GRB 220124A
DATE: 22/01/25 15:07:51 GMT
FROM: Dmitry Svinkin at Ioffe Institute <svinkin@mail.ioffe.ru>

A.S. Kezyrev, D.V. Golovin, M.I. Litvak, T.G. Mitrofanov, and A.B. on behalf of the HEND/Mars Odyssey team,

D. Svinkin, D. Frederiks, A. Ridaia, A. Lyzenko, and T. Cline on behalf of the Konus-Wind team,

A. von Kienlin, X. Zhang, A. Rau, V. Savchenko, E. Bozza, and C. Ferrigno, on behalf of the INTEGRAL SPI-ACS GRB team,

S. Barthelay, J. Cummings, H. Krims, D. Palmer, and A. Tohuvavohu on behalf of the Swift-BAT team, and

W. Boynton, C. Fellows, K. Harshman, H. Enos, R. Starr, and A.S. Gardner on behalf of the GRS-Odyssey GRB team, report:

The long-duration GRB 220124A (AstroSat-CZTI detection: Prasad et al., GCN Circ. 31521; GECAM detection: Liu et al., GCN Circ. 31524) has been detected by AstroSat (CZTI), Konus-Wind, Swift (BAT), INTEGRAL (SPI-ACS), Mars-Odyssey (HEND), CALET (CGRM), and GECAM-B (GRD), so far, at about 13958 s UT (03:37:38). The burst was outside the coded field of view of the BAT.

We have triangulated it to a preliminary 95% error box whose coordinates are:

RA(2000), deg	Dec(2000), deg
Center:	
241.945 (16h 07m 47s)	+41.979 (+41d 58' 45")
Corners:	
244.220 (16h 16m 53s)	+42.801 (+42d 48' 62")
239.771 (15h 59m 05s)	+41.164 (+41d 09' 51")
239.711 (15h 58m 51s)	+41.602 (+41d 04' 54")
244.155 (16h 16m 37s)	+42.723 (+42d 43' 23")

The error box area is 678 sq. arcmin, and its maximum dimension is 3.8 deg (the minimum one is 2.5 arcmin). The Sun's distance was 84 deg.

02/08/2022

Example Walk-Thru

- Search for galaxies in this polygon in SIMBAD:

Portal Simbad Vizier Aladin X-Match Other Help

SIMBAD: Query by criteria

other query Identifier query Coordinate query **Criteria query** Reference query Basic query Script submission TAP Output options Help

Enter a search expression:

region(polygon ICRS, 244.220 +42.801 , 239.771 +41.164 , 239.711 +41.082 ,244.155 +42.723) & Otype = 'G'

Criteria queries may require some time, especially if they are complex or involve a large number of objects. Limited to 5 criteria. Please, wait for their completion if it is the case.

Example:
 $ra > 15 \& ra < 30 \& dec > 70$
 $(cat = 'PPM' | cat = 'HIP')$
 $Rot.vsinl > 10.0$

Return:
 object count
 display maximum 10000 objects
 get references from the selected objects

Enter the name of an ASCII file containing a search expression:

Browse... No file selected.

submit file clear

Example Walk-Thru: Simbad result:

region(polygon ICRS, 244.220 +42.801 , 239.771 +41.164 , 239.711 +41.082 ,244.155 +42.723) & Otype = 'G'


other query Identifier query Coordinate query **Criteria query** Reference query Basic query Script submission TAP Output options Help

Query : region(polygon ICRS, 244.220 +42.801 , 239.771 +41.164 , 239.711 +41.082 ,244.155 +42.723) & Otype = 'G'

Number of rows : 9

Show 100 entries

N°	Identifier	Otype	ICRS (J2000) RA	ICRS (J2000) DEC	Mag U
1	LEDA 56789	G	16 02 20.405	+41 27 39.75	
2	SDSS J160924.76+420624.8	G	16 09 24.764	+42 06 24.86	
3	LEDA 2205680	G	16 16 40.224	+42 45 51.03	
4	LEDA 2176874	G	15 59 43.2	+41 10 32	
5	2MASX J16091874+4207067	G	16 09 18.7434001848	+42 07 06.901291056	
6	2MASX J16145890+4237371	G	16 14 58.916	+42 37 37.22	
7	2MASX J16094346+4207507	G	16 09 43.495	+42 07 50.68	
8	SDSS J161023.56+421338.6	G	16 10 23.564	+42 13 38.68	
9	SDSS J155915.34+410729.1	G	15 59 15.348	+41 07 29.11	



AAVSO Alerts and Campaigns, HEN Forum


- Visit us at aavso.org

- High Energy Network Section forum:

<https://www.aavso.org/forum/8220>

- Always worth to keep an eye on current alerts and campaigns

<https://www.aavso.org/observers>



The AAVSO International High Energy Network Observing Section Webinar “*How to Observe Optical Counterparts of High Energy Astronomical Transients*”, presented by section co-lead Heinz-Bernd Eggenstein, can be found here. [How to Observe Optical Counterparts of High Energy Astronomical Transients](#).

Acknowledgements

Contributors to this page are Heinz-Bernd Eggenstein, Arto Oksanen, and Filipp Romanov.

The GRB observational techniques presented herein were developed and successfully utilized by AAVSO members Arto Oksanen and Filipp Romanov. Their work is critical to further research in GRB astrophysics.

References (with links)

[New Supernova Alert System Promises Early Access to Spectacles in Space \(Scientific American; February 22, 2021\)](#)

SNEWS (SuperNova Early Warning System); Pietro Antonioli *et al* 2004 *New J. Phys.* 6 114; [SNEWS: the SuperNova Early Warning System](#)

LIGO (Laser Interferometer Gravitational-wave Observatory); The LIGO Scientific Collaboration *et al* 2021 [Search for anisotropic gravitational-wave backgrounds using data from Advanced LIGO's and Advanced Virgo's first three observing runs](#)

VIRGO; B.P. Abbott *et al* (LIGO Scientific Collaboration and Virgo Collaboration) *Phys. Rev. Lett.* 116, 061102 2016 [Observation of Gravitational Waves from a Binary Black Hole Merger](#)

Personal communications among the contributors to this page; Heinz-Bernd Eggenstein, Arto Oksanen, and Filipp Romanov.

Oksanen, A., *et al.* 2008, "Discovery and Observations of the Optical Afterglow of GRB 071010B", *JAAVSO* Volume 36, 2008.

Oksanen, A., *et al.* 2002, "Gamma-Ray Burst Optical Afterglow Observations at Nyrola Observatory", *JAAVSO* Volume 30, 2002.

AAVSO High Energy Network Observing Section webinar (July 25, 2020): https://www.youtube.com/watch?v=mfr0ZoPh2_o&list=PLnZ_rvnR35rf3rDie-XWhapGZlqYlrJwn&index=5 (Observing GRB Optical Afterglows-Arto Oksanen).